

ON CERTAIN CUNTZ-PIMSNER ALGEBRAS

ALEX KUMJIAN

ABSTRACT. Let A be a separable unital C^* -algebra and let $\pi : A \rightarrow \mathcal{L}(\mathfrak{H})$ be a faithful representation of A on a separable Hilbert space $\mathfrak{H}$ such that $\pi(A) \cap \mathcal{K}(\mathfrak{H}) = \{0\}$. We show that $\mathcal{O}_E$, the Cuntz-Pimsner algebra associated to the Hilbert A -bimodule $E = \mathfrak{H} \otimes_C A$, is simple and purely infinite. If A is nuclear and belongs to the bootstrap class to which the UCT applies, then the same applies to $\mathcal{O}_E$. Hence by the Kirchberg-Phillips Theorem the isomorphism class of $\mathcal{O}_E$ only depends on the K -theory of A and the class of the unit.

In his seminal paper [Pm], Pimsner constructed a C^* -algebra $\mathcal{O}_E$ from a Hilbert bimodule over a C^* -algebra A as a quotient of a concrete C^* -algebra $\mathcal{T}_E$, an analogue of the Toeplitz algebra, acting on the Fock space associated to E . There has recently been much interest in these Cuntz-Pimsner algebras (or Cuntz-Krieger-Pimsner algebras), which generalize both crossed products by $\mathbb{Z}$ and Cuntz-Krieger algebras, as well as the associated Toeplitz algebras. The structure of these C^* -algebras is not yet fully understood, although considerable progress has been made. For example, Pimsner found a six-term exact sequence for the K -theory of $\mathcal{O}_E$ which generalizes the Pimsner-Voiculescu exact sequence (see [Pm, Theorem 4.8]); conditions for simplicity were found in [Sc2, MS, KPW1, DPW] and for pure infiniteness in [Z].

The purpose of the present note is to analyze the structure of Cuntz-Pimsner algebras associated to a certain class of Hilbert bimodules. Let A be a separable unital C^* -algebra and let $\pi : A \rightarrow \mathcal{L}(\mathfrak{H})$ be a faithful representation of A on a separable Hilbert space $\mathfrak{H}$ such that $\pi(A) \cap \mathcal{K}(\mathfrak{H}) = \{0\}$. Then $E = \mathfrak{H} \otimes_C A$ is a Hilbert bimodule over A in a natural way. We show that $\mathcal{O}_E$ is separable, simple and purely infinite. If A is nuclear and in the bootstrap class, then the same holds for $\mathcal{O}_E$ and thus by the Kirchberg-Phillips theorem the isomorphism class of $\mathcal{O}_E$ is completely determined by the K -theory of A together with the class of the unit (since $\mathcal{O}_E$ is KK -equivalent to A).

Many examples of Cuntz-Pimsner algebras found in the literature arise from Hilbert bimodules which are finitely generated and projective; in such cases the left action must consist entirely of compact operators. Our examples do not fall in this class; in fact, the left action has trivial intersection with the compacts. And this has some interesting consequences: $\mathcal{O}_E \cong \mathcal{T}_E$ (see [Pm, Corollary 3.14]) and the natural embedding $A \hookrightarrow \mathcal{O}_E$ induces a KK -equivalence (see [Pm, Corollary 4.5]).

In §1 we review some basic facts concerning the construction of $\mathcal{T}_E$ as operators on the Fock space of E and the gauge action $\lambda : \mathbb{T} \rightarrow \text{Aut}(\mathcal{T}_E)$. We assume that the left action of A does not meet the compacts $\mathcal{K}(E)$ and identify $\mathcal{O}_E$ with $\mathcal{T}_E$. The fixed point algebra $\mathcal{F}_E$, the analogue of the AF-core of a Cuntz-Krieger algebra, contains a canonical descending sequence of essential ideals indexed by $\mathbb{N}$ with trivial intersection. The crossed product $\mathcal{O}_E \rtimes_\lambda \mathbb{T}$ has a similar collection of essential ideals indexed by $\mathbb{Z}$ on which the dual group of automorphisms acts in a natural way. By Takesaki-Takai duality

$$\mathcal{O}_E \otimes \mathcal{K}(L^2(\mathbb{T})) \cong (\mathcal{O}_E \rtimes_\lambda \mathbb{T}) \rtimes_{\hat{\lambda}} \mathbb{Z};$$

Date: 28 August 2001.

1991 Mathematics Subject Classification. Primary 46L05; Secondary 46L55.

Key words and phrases. C^* -algebra, Hilbert bimodule, simple, purely infinite.

Research partially supported by NSF grant DMS-9706982.

hence, much of the structure of $\mathcal{O}_E$ is revealed through an analysis of the double crossed product. In §2 we show that if E is the Hilbert bimodule over A associated to a representation as described above, then for every nonzero positive element $d \in \mathcal{O}_E$ there is a $z \in \mathcal{O}_E$ so that $z^*dz = 1$; it follows that $\mathcal{O}_E$ is simple and purely infinite (see Theorem 2.8). The proof of this proceeds through a sequence of lemmas and is patterned on the proof of [Rø, Theorem 2.1], which is in turn based on a key lemma of Kishimoto (see [Ks, Lemma 3.2]). Our argument uses the version of this lemma found in [OP3, Lemma 7.1] and this requires that we show that the Connes spectrum of the dual action is full (this is also an ingredient in the proof of simplicity found in [DPW]). We invoke a version of a key lemma of Rørdam for crossed products by $\mathbb{Z}$ which arise from automorphisms with full Connes spectrum. The fact that $\mathcal{O}_E$ embeds equivariantly into $(\mathcal{O}_E \rtimes_{\lambda} \mathbb{T}) \rtimes_{\hat{\lambda}} \mathbb{Z}$ allows us to apply this lemma to $\mathcal{O}_E$. In §3 we use the Kirchberg-Phillips theorem to collect some consequences of this theorem as indicated above and discuss certain connections with reduced (amalgamated) free products.

We fix some notation and terminology. Given a C^* -algebra B we let $\hat{B}$ denote its spectrum, that is, the collection of irreducible representations modulo unitary equivalence endowed with the Jacobson topology (see [Pd, §4.1]). If I is an ideal in a C^* -algebra B , then every irreducible representation of I extends uniquely to an irreducible representation of B . This allows one to identify $\hat{I}$ with an open subset of $\hat{B}$, the complement of which consists of the classes of irreducible representations which vanish on I . Given a $*$ -automorphism β of a C^* -algebra B , let $\Gamma(\beta)$ denote the Connes spectrum of β (see [O, Co] or [Pd, §8.8]); recall that

$$\Gamma(\beta) = \bigcap_H \text{Sp}(\beta|_H)$$

where the intersection is taken over all β -invariant hereditary subalgebras H . A C^* -algebra is said to be purely infinite if every hereditary subalgebra contains an infinite projection.

I wish to thank D. Shlyakhtenko for certain helpful remarks relating to material in §3.

1. PRELIMINARIES

We review some basic facts concerning Cuntz-Pimsner algebras; we shall be mainly interested in those which arise from bimodules for which the left action has trivial intersection with the compacts (see Remark 1.3). Let A be a C^* -algebra.

Definition 1.1. (see [Ri, Ka, L]) *Let E be a right A -module. Then E is said to be a (right) pre-Hilbert A -module if it is equipped with an A -valued inner product $\langle \cdot, \cdot \rangle_A$ which satisfies the following conditions for all $\xi, \eta, \zeta \in E$, $s, t \in \mathbb{C}$, and $a \in A$:*

- i. $\langle \xi, s\eta + t\zeta \rangle_A = s\langle \xi, \eta \rangle_A + t\langle \xi, \zeta \rangle_A$
- ii. $\langle \xi, \eta a \rangle_A = \langle \xi, \eta \rangle_A a$
- iii. $\langle \eta, \xi \rangle_A = \langle \xi, \eta \rangle_A^*$
- iv. $\langle \xi, \xi \rangle_A \geq 0$ and $\langle \xi, \xi \rangle_A = 0$ only if $\xi = 0$.

E is said to be a (right) Hilbert A -module if it is complete in the norm: $\|\xi\| = \|\langle \xi, \xi \rangle_A\|^{1/2}$.

Let E be a Hilbert A -module. Then E is said to be *full* if the span of the values of the inner product is dense. The collection of bounded adjointable operators on E , $\mathcal{L}(E)$, is a C^* -algebra. The closure of the span of operators of the form $\theta_{\xi, \eta}$ for $\xi, \eta \in E$ (where $\theta_{\xi, \eta}(\zeta) = \xi \langle \eta, \zeta \rangle_A$ for $\zeta \in E$) forms an essential ideal in $\mathcal{L}(E)$ which is denoted $\mathcal{K}(E)$. A Hilbert space is a Hilbert module over $\mathbb{C}$.

Definition 1.2. *Let E be a Hilbert A -module and $\varphi : A \rightarrow \mathcal{L}(E)$ be an injective $*$ -homomorphism. Then the pair (E, φ) is said to be Hilbert bimodule over A (or Hilbert A -bimodule).*

Pimsner defines the Cuntz-Pimsner algebra $\mathcal{O}_E$ as a quotient of the analogue of the Toeplitz algebra, $\mathcal{T}_E$, generated by creation operators on the Fock space of E (see [Pm]). The injectivity

of φ is not really necessary (see [Pm, Remark 1.2.1]). We will henceforth assume that E is full (see [Pm, Remark 1.2.3]).

The Fock space of E is the Hilbert A -module

$$\mathcal{E}_+ = \bigoplus_{n=0}^{\infty} E^{\otimes n}$$

where $E^{\otimes 0} = A$ and for $n > 0$, $E^{\otimes n}$ is the n -fold tensor product:

$$E^{\otimes n} = E \otimes_A \cdots \otimes_A E.$$

Note that $\mathcal{E}_+$ is also a Hilbert A -bimodule with left action defined by $\varphi_+(a)b = ab$ for $a, b \in A = E^{\otimes 0}$ and

$$\varphi_+(a)(\xi_1 \otimes \cdots \otimes \xi_n) = \varphi(a)\xi_1 \otimes \cdots \otimes \xi_n$$

for $a \in A$ and $\xi_1 \otimes \cdots \otimes \xi_n \in E^{\otimes n}$.

Then $\mathcal{T}_E \subset \mathcal{L}(\mathcal{E}_+)$ is the C^* -algebra generated by the creation operators T_ξ for $\xi \in E$ where $T_\xi(a) = \xi a$ and

$$T_\xi(\xi_1 \otimes \cdots \otimes \xi_n) = \xi \otimes \xi_1 \otimes \cdots \otimes \xi_n.$$

Observe that $T_\xi^* T_\eta = \varphi_+(\langle \xi, \eta \rangle_A)$ for $\xi, \eta \in E$. Since E is full, $\varphi_+(A) \subset \mathcal{T}_E$; let $\iota : A \hookrightarrow \mathcal{T}_E$ denote the embedding. Note that one may define T_ξ for $\xi \in E^{\otimes n}$ in an analogous manner and that we have $T_\xi^* T_\eta = \iota(\langle \xi, \eta \rangle_A)$ for $\xi, \eta \in E^{\otimes n}$.

There is an embedding $\iota_n : \mathcal{K}(E^{\otimes n}) \hookrightarrow \mathcal{T}_E$ (identify $\mathcal{K}(E^{\otimes 0})$ with A), given for $n > 0$ by $\iota_n(\theta_{\xi, \eta}) = T_\xi T_\eta^*$ for $\xi, \eta \in E^{\otimes n}$. Note that such operators preserve the grading of $\mathcal{E}_+$ and that there is an embedding $\mathcal{K}(E^{\otimes n}) \hookrightarrow \mathcal{L}(E^{\otimes m})$ for $m \geq n$. Let C_n denote the C^* -subalgebra of $\mathcal{T}_E$ generated by operators of the form $T_\xi T_\eta^*$ for $\xi, \eta \in E^{\otimes k}$ with $k \leq n$ (by convention $C_0 = \iota(A)$). Then the C_n form an ascending family of C^* -subalgebras.

Remark 1.3. Suppose $\varphi(A) \cap \mathcal{K}(E) = \{0\}$; then the natural map $C_n \rightarrow \mathcal{L}(E^{\otimes m})$ is an embedding for $m \geq n$. By [Pm, Corollary 3.14] $\mathcal{T}_E \cong \mathcal{O}_E$ and the inclusion $A \hookrightarrow \mathcal{O}_E$ induces a KK -equivalence (see [Pm, Corollary 4.5]). Under the isomorphism of $\mathcal{T}_E$ with $\mathcal{O}_E$, $\overline{\cup_n C_n}$ is mapped to $\mathcal{F}_E$, the analog of the AF core of a Cuntz-Krieger algebra.

For the remainder of this section we shall tacitly assume that $\varphi(A) \cap \mathcal{K}(E) = \{0\}$ and identify $\mathcal{T}_E$ with $\mathcal{O}_E$.

Proposition 1.4. *For each $n \in \mathbb{N}$ the C^* -subalgebra, J_n , generated by $\iota_k(\mathcal{K}(E^{\otimes k}))$ for $k \geq n$ is an essential ideal in $\mathcal{F}_E$. We obtain a descending sequence of ideals*

$$J_0 \supset J_1 \supset J_2 \supset \cdots$$

with $J_0 = \mathcal{F}_E$ and $\cap_n J_n = \{0\}$. Furthermore, $J_n/J_{n+1} \cong \mathcal{K}(E^{\otimes n})$ (thus J_n/J_{n+1} is strong Morita equivalent to A) and the restriction of the quotient map yields an isomorphism $C_n \cong \mathcal{F}_E/J_{n+1}$.

Proof. Given $n \in \mathbb{N}$ it is clear that J_n is an ideal (see [Pm, Definition 2.1]). To see that J_n is essential it suffices to show that for every m and nonzero element $c \in C_m$ there is an element $d \in \mathcal{K}(E^{\otimes k})$ for some $k \geq n$ such that $c\iota_k(d) \neq 0$. Let k be an integer with $k \geq \max(m, n)$; since the map from C_m to $\mathcal{L}(E^{\otimes k})$ is an embedding for $k \geq m$, $c\xi \neq 0$ for some $\xi \in E^{\otimes k}$. Then $cT_\xi T_\xi^* \neq 0$ and we take $d = \theta_{\xi, \xi}$.

The J_n form a descending sequence of ideals by construction. Since $\varphi(A) \cap \mathcal{K}(E) = \{0\}$, $C_m \cap J_n = \{0\}$ for $m < n$. Hence, $\cap_n J_n = \{0\}$, for $\mathcal{F}_E$ is the inductive limit of the C_m . Further, for each n we have

$$J_n = \iota_n(\mathcal{K}(E^{\otimes n})) + J_{n+1} \quad \text{and} \quad \iota_n(\mathcal{K}(E^{\otimes n})) \cap J_{n+1} = \{0\};$$

it follows that $J_n/J_{n+1} \cong \mathcal{K}(E^{\otimes n})$. Finally, since

$$\mathcal{F}_E = C_n + J_{n+1} \quad \text{and} \quad C_n \cap J_{n+1} = \{0\},$$

we have $C_n \cong \mathcal{F}_E/J_{n+1}$. □

There is a strongly continuous action

$$\lambda : \mathbb{T} \rightarrow \text{Aut}(\mathcal{O}_E)$$

such that $\lambda_t(T_\xi) = tT_\xi$. The fixed point algebra under this action is $\mathcal{F}_E$ and we have a faithful conditional expectation $P_E : \mathcal{O}_E \rightarrow \mathcal{F}_E$ given by

$$P_E(x) = \int_{\mathbb{T}} \lambda_t(x) dt.$$

Consider the spectral subspaces of $\mathcal{O}_E$ under this action: for $n \in \mathbb{Z}$

$$(\mathcal{O}_E)_n = \{x \in \mathcal{O}_E : \lambda_t(x) = t^n x \text{ for all } t \in \mathbb{T}\}.$$

Remark 1.5. Note that $(\mathcal{O}_E)_n$ is the closure of the span of elements of the form $T_\xi T_\eta^*$ where $\xi \in E^{\otimes k}$ and $\eta \in E^{\otimes l}$ with $n = k - l$. For $n \geq 0$ and $x \in (\mathcal{O}_E)_n$ we have $x^*x \in \mathcal{F}_E$ and $xx^* \in J_n$. We may regard $(\mathcal{O}_E)_n$ as an J_n - $\mathcal{F}_E$ -equivalence bimodule (see [Ri]); hence, J_n is strong Morita equivalent to $\mathcal{F}_E$ for each $n \geq 0$. If we regard $(\mathcal{O}_E)_1$ as a Hilbert $\mathcal{F}_E$ -bimodule we have (cf. [Pm, §2] and [Sc2, §1.4])

$$(\mathcal{O}_E)_1 \cong E \otimes_A \mathcal{F}_E,$$

where the isomorphism is implemented by the map $\xi \otimes a \mapsto T_\xi a$. The crossed product $\mathcal{O}_E \rtimes_\lambda \mathbb{T}$ may be identified with the closure of the subalgebra of $\mathcal{O}_E \otimes \mathcal{K}(\ell^2(\mathbb{Z}))$ consisting of finite sums of the form

$$\sum x_{ij} \otimes e_{ij}$$

where e_{ij} are the standard rank one partial isometries in $\mathcal{K}(\ell^2(\mathbb{Z}))$ and $x_{ij} \in (\mathcal{O}_E)_{j-i}$.

Let $\widehat{\lambda} : \mathbb{Z} \rightarrow \text{Aut}(\mathcal{O}_E \rtimes_\lambda \mathbb{T})$ denote the dual automorphism group.

Proposition 1.6. *There is an embedding $\epsilon : \mathcal{F}_E \hookrightarrow \mathcal{O}_E \rtimes_\lambda \mathbb{T}$ onto a corner and a collection of essential ideals $\{I_n\}_{n \in \mathbb{Z}}$ in $\mathcal{O}_E \rtimes_\lambda \mathbb{T}$ satisfying the following conditions:*

- i. *For all $n \in \mathbb{Z}$, $\mathcal{F}_E$ is strong Morita equivalent to I_n and A is strong Morita equivalent to I_n/I_{n+1} .*
- ii. *For all $n \geq 0$, $\epsilon(J_n) = \epsilon(1)I_n\epsilon(1)$.*
- iii. *$I_n \subset I_m$ if $m \leq n$.*
- iv. *$\bigcap_n I_n = \{0\}$*
- v. *$\bigcup_n I_n = \mathcal{O}_E \rtimes_\lambda \mathbb{T}$*
- vi. *$\widehat{\lambda}_k(I_n) = I_{n+k}$*

Proof. We use the identification of $\mathcal{O}_E \rtimes_\lambda \mathbb{T}$ with a C^* -subalgebra of $\mathcal{O}_E \otimes \mathcal{K}(\ell^2(\mathbb{Z}))$ given in Remark 1.5. For each n let I_n be the ideal generated by $p_n = 1 \otimes e_{nn}$. Since $\mathcal{F}_E = (\mathcal{O}_E)_0$, it follows that $\mathcal{F}_E$ is isomorphic to the corner determined by p_n and thus is strong Morita equivalent to I_n . The desired embedding $\epsilon : \mathcal{F}_E \hookrightarrow \mathcal{O}_E \rtimes_\lambda \mathbb{T}$ is given by $\epsilon(a) = a \otimes e_{00}$. Given an element of the form $a_{mn} = x_{mn} \otimes e_{mn}$ in $\mathcal{O}_E \rtimes_\lambda \mathbb{T}$ with $m \leq n$, we have

$$a_{mn}^* a_{mn} = x_{mn}^* x_{mn} \otimes e_{nn} \quad \text{and} \quad a_{mn} a_{mn}^* = x_{mn} x_{mn}^* \otimes e_{mm}$$

with $x_{mn} x_{mn}^* \in J_{n-m}$; since p_n may be expressed as a finite sum of elements of the form $a_{mn}^* a_{mn}$, it follows that $I_n \subset I_m$ and that $p_m I_n p_m = J_{n-m} \otimes e_{mm}$. Thus $\epsilon(J_n) = \epsilon(1)I_n\epsilon(1)$ for all $n \geq 0$. Assertion (vi) follows from the fact that $\widehat{\lambda}_k(p_n) = 1 \otimes p_{n+k}$. The remaining assertions follow from Proposition 1.4. $\square$

2. $\mathcal{O}_E$ IS SIMPLE AND PURELY INFINITE

Let A be a separable unital C^* -algebra and let $\pi : A \rightarrow \mathcal{L}(\mathfrak{H})$ be a faithful representation of A on a separable Hilbert space $\mathfrak{H}$.

Proposition 2.1. *With A and $\pi : A \rightarrow \mathcal{L}(\mathfrak{H})$ as above,*

$$E = \mathfrak{H} \otimes_{\mathbb{C}} A$$

is a full Hilbert bimodule over A under the operations

$$\langle \xi \otimes a, \eta \otimes b \rangle_A = \langle \xi, \eta \rangle a^* b, \quad \varphi(a)(\xi \otimes b) = \pi(a)\xi \otimes b$$

for all $\xi, \eta \in \mathfrak{H}$ and $a, b \in A$. Moreover, if $\pi(A) \cap \mathcal{K}(\mathfrak{H}) = \{0\}$, then $\varphi(A) \cap \mathcal{K}(E) = \{0\}$ and $\mathcal{O}_E \cong \mathcal{T}_E$.

Proof. Note that $E = \mathfrak{H} \otimes_{\mathbb{C}} A$ is the tensor product of the Hilbert A - $\mathbb{C}$ -bimodule $\mathfrak{H}$ and the Hilbert $\mathbb{C}$ - A -bimodule A as defined by Rieffel in [Ri] (see also [L, Ch. 4]). The natural map from $\mathcal{L}(\mathfrak{H})$ to $\mathcal{L}(E) = \mathcal{L}(\mathfrak{H} \otimes_{\mathbb{C}} A)$ induces an embedding $\mathcal{L}(\mathfrak{H})/\mathcal{K}(\mathfrak{H}) \hookrightarrow \mathcal{L}(E)/\mathcal{K}(E)$ (since $\mathcal{K}(\mathfrak{H})$ is mapped into $\mathcal{K}(E)$ and the Calkin algebra $\mathcal{L}(\mathfrak{H})/\mathcal{K}(\mathfrak{H})$ is simple). Hence, if $\pi(A) \cap \mathcal{K}(\mathfrak{H}) = \{0\}$, then $\varphi(A) \cap \mathcal{K}(E) = \{0\}$. The last assertion, $\mathcal{O}_E \cong \mathcal{T}_E$, follows by [Pm, Corollary 3.14]. $\square$

Henceforth, we assume that $\pi(A) \cap \mathcal{K}(\mathfrak{H}) = \{0\}$ and identify $\mathcal{O}_E$ with $\mathcal{T}_E$. The aim of this section is to show that $\mathcal{O}_E$ is simple and purely infinite. Simplicity may be proven directly by invoking [Sc2, Theorem 3.9]: if A is unital and E is full, then $\mathcal{O}_E$ is simple if and only if E is minimal and nonperiodic. Lemma 2.3 would then be a consequence of [OP1, Theorem 6.5]. We follow a more indirect route patterned on the proof of [Rø, Theorem 2.1]; this will also show that $\mathcal{O}_E$ is purely infinite.

Remark 2.2. With $E = \mathfrak{H} \otimes_{\mathbb{C}} A$ as above, we have $E^{\otimes n} \cong \mathfrak{H}^{\otimes n} \otimes_{\mathbb{C}} A$ via the map

$$(\xi_1 \otimes a_1) \otimes (\xi_2 \otimes a_2) \otimes \cdots \otimes (\xi_n \otimes a_n) \mapsto (\xi_1 \otimes \pi(a_1)\xi_2 \otimes \cdots \otimes \pi(a_{n-1})\xi_n) \otimes a_n;$$

similarly, if $\sigma : A \rightarrow \mathcal{L}(\mathfrak{K})$ is a representation of A on a Hilbert space $\mathfrak{K}$, then

$$E^{\otimes n} \otimes_A \mathfrak{K} \cong E^{\otimes n-1} \otimes_A \mathfrak{H} \otimes \mathfrak{K}.$$

Recall that the action of $\mathcal{F}_E$ on Fock space preserves the natural grading. Let $\tilde{\sigma}_n$ denote the representation of $\mathcal{F}_E$ on $E^{\otimes n} \otimes_A \mathfrak{K}$ given by left action on $E^{\otimes n}$. Then the restriction of $\tilde{\sigma}_n$ to C_{n-1} is faithful: indeed, this follows from the facts that the natural map

$$\mathcal{L}(E^{\otimes n-1}) \rightarrow \mathcal{L}(E^{\otimes n-1} \otimes_A \mathfrak{H} \otimes \mathfrak{K}) \cong \mathcal{L}(E^{\otimes n} \otimes_A \mathfrak{K})$$

is an embedding (since π is faithful) and that $\tilde{\sigma}_n|_{\mathcal{K}(E^{\otimes n-1})}$ factors through $\mathcal{L}(E^{\otimes n-1})$. Note that $\tilde{\sigma}_n$ is equivalent to the representation of $\mathcal{F}_E$ obtained from σ as follows: use the strong Morita equivalence between A and J_n/J_{n+1} to obtain a representation of J_n/J_{n+1} and extend this to a representation of $\mathcal{F}_E$. Since the restriction of $\tilde{\sigma}_n$ to C_{n-1} is faithful, $\ker \tilde{\sigma}_n \subset J_n$ (see Proposition 1.4). It follows that the closure of a point in $\widehat{J_n} - \widehat{J_{n+1}}$ contains the complement of $\widehat{J_n}$. A similar assertion holds for $\mathcal{O}_E \rtimes_{\lambda} \mathbb{T}$: for any $n \in \mathbb{Z}$ the closure of a point in $\widehat{I_n} - \widehat{I_{n+1}}$ contains the complement of $\widehat{I_n}$.

Lemma 2.3. *With A and E as above, $\Gamma(\widehat{\lambda}_1) = \mathbb{T}$ where $\widehat{\lambda}$ is the dual action of $\mathbb{Z}$ on $\mathcal{O}_E \rtimes_{\lambda} \mathbb{T}$.*

Proof. By [OP2, Theorem 4.6] it suffices to find a dense invariant subset of $(\mathcal{O}_E \rtimes_{\lambda} \mathbb{T})^{\wedge}$ on which $\widehat{\lambda}_1^*$ acts freely. That is, we must find an irreducible representation σ of $\mathcal{O}_E \rtimes_{\lambda} \mathbb{T}$ such that, $\{[\sigma \circ \widehat{\lambda}_n] : n \in \mathbb{Z}\}$, the orbit of the unitary equivalence class of σ under $\widehat{\lambda}^*$, is dense in $(\mathcal{O}_E \rtimes_{\lambda} \mathbb{T})^{\wedge}$ and $[\sigma \circ \widehat{\lambda}_m] \neq [\sigma \circ \widehat{\lambda}_n]$ if $m \neq n$. Let σ_0 be an irreducible representation of A and use the strong Morita equivalence between A and I_0/I_1 to obtain an irreducible representation σ' of I_0/I_1 . Then

σ , the extension of σ' to $\mathcal{O}_E \rtimes_\lambda \mathbb{T}$, is also irreducible. The classes $[\sigma \circ \widehat{\lambda}_n]$ are distinct, for if $m < n$, $\sigma \circ \widehat{\lambda}_m$ vanishes on I_n . Moreover, for each $n \in \mathbb{Z}$ the closure of $[\sigma \circ \widehat{\lambda}_n]$ in $(\mathcal{O}_E \rtimes_\lambda \mathbb{T})^\wedge$ includes the classes of all irreducible representations which vanish on I_n (since $[\sigma \circ \widehat{\lambda}_n] \in \widehat{I_n} - \widehat{I_{n+1}}$, see Remark 2.2). Hence, $\{[\sigma \circ \widehat{\lambda}_n] : n \in \mathbb{Z}\}$ is dense in $(\mathcal{O}_E \rtimes_\lambda \mathbb{T})^\wedge$. $\square$

Using Takesaki-Takai duality we show below that a C^* -algebra D equipped with an action α of $\mathbb{T}$ may be embedded equivariantly as a corner in $(D \rtimes_\alpha \mathbb{T}) \rtimes_{\widehat{\alpha}} \mathbb{Z}$. This fact is related to Rosenberg's observation that the fixed point algebra under a compact group action embeds as a corner in the crossed product (see [Ro]).

Proposition 2.4. *Given a unital C^* -algebra D and a strongly continuous action $\alpha : \mathbb{T} \rightarrow \text{Aut}(D)$, there is an isomorphism ψ of D onto a full corner of $(D \rtimes_\alpha \mathbb{T}) \rtimes_{\widehat{\alpha}} \mathbb{Z}$ which is equivariant in the sense that $\widehat{\alpha}_t \circ \psi = \psi \circ \alpha_t$ for all $t \in \mathbb{T}$. Moreover, $\psi(1) \in D \rtimes_\alpha \mathbb{T}$.*

Proof. By Takesaki-Takai duality [Pd, 7.9.3] there is an isomorphism

$$\gamma : D \otimes \mathcal{K}(L^2(\mathbb{T})) \cong (D \rtimes_\alpha \mathbb{T}) \rtimes_{\widehat{\alpha}} \mathbb{Z},$$

which is equivariant with respect to $\alpha \otimes \text{Ad } \rho$ and $\widehat{\alpha}$ (where ρ is the right regular representation of $\mathbb{T}$ on $L^2(\mathbb{T})$). The desired embedding is obtained by finding an $\text{Ad } \rho$ invariant minimal projection p in $\mathcal{K}(L^2(\mathbb{T}))$ (cf. [Ro]): set $\psi(d) = \gamma(d \otimes p)$ for $d \in D$. Since ψ is equivariant, $\psi(1)$ is in the fixed point algebra of $\widehat{\alpha}$; hence, $\psi(1) \in D \rtimes_\alpha \mathbb{T}$. $\square$

The following lemma is adapted from [Rø, Lemma 2.4]; the proof is patterned on Rørdam's but we substitute [OP3, Lemma 7.1] for [Ks, Lemma 3.2].

Lemma 2.5. *Let B be a C^* -algebra and let β be an automorphism of B such that $\Gamma(\beta) = \mathbb{T}$ and let P denote the canonical conditional expectation from $B \rtimes_\beta \mathbb{Z}$ to B . Then for every positive element $y \in B \rtimes_\beta \mathbb{Z}$ and $\varepsilon > 0$ there are positive elements $x, b \in B$ such that*

$$\|b\| > \|P(y)\| - \varepsilon, \quad \|x\| \leq 1 \quad \text{and} \quad \|xyx - b\| < \varepsilon.$$

If y is in the corner determined by a projection $p \in B$, then x, b may also be chosen to be in the corner.

Proof. As in the proof of [Rø, Lemma 2.4] we may assume (by perturbing y if necessary) that y is of the form

$$y = y_{-n}u^{-n} + \cdots + y_{-1}u^{-1} + y_0 + y_1u + \cdots + y_nu^n$$

for some n where $y_j \in B$ and u is the canonical unitary in $B \rtimes_\beta \mathbb{Z}$ implementing the automorphism β ; note that $y_0 = P(y)$ is positive.

By [OP3, Theorem 10.4] β^k is properly outer for all $k \neq 0$. Hence, by [OP3, Lemma 7.1] there is a positive element x with $\|x\| = 1$ such that

$$\|xy_0x\| > \|y_0\| - \varepsilon, \quad \text{and} \quad \|xy_ku^kx\| = \|xy_k\beta^k(x)\| < \varepsilon/2n$$

for $0 < |k| \leq n$. Set $b = xy_0x$; then a straightforward calculation yields $\|xyx - b\| < \varepsilon$. We now verify the last assertion. Suppose that y is in the corner determined by a projection $p \in B$; we may again assume that y is of the above form. Since P is a conditional expectation onto B , $y_0 = P(y)$ is also in the corner determined by p . In the proof of [OP3, Lemma 7.1] the positive element x is constructed in the hereditary subalgebra determined by y_0 ; hence we may assume that x and therefore also $b = xy_0x$ lies in the same corner. $\square$

Recall that C_n is the C^* -subalgebra of $\mathcal{F}_E$ generated by operators of the form $T_\xi T_\eta^*$ for $\xi, \eta \in E^{\otimes k}$ with $k \leq n$ and that they form an ascending family of C^* -subalgebras with dense union. The subspace $E^{\otimes n}$ is left invariant by C_n and one has an embedding $C_n \hookrightarrow \mathcal{L}(E^{\otimes n})$.

Lemma 2.6. *Given a positive element $c \in C_n$ and $\varepsilon > 0$, there is $\xi \in E^{\otimes n}$ with $\|\xi\| = 1$ such that $T_\xi^* c T_\xi \in C_0$ and $\|T_\xi^* c T_\xi\| > \|c\| - \varepsilon$.*

Proof. The first assertion follows from a straightforward calculation: given $c \in C_n$ and $\xi \in E^{\otimes n}$, then $c\xi \in E^{\otimes n}$ and

$$T_\xi^* c T_\xi = T_\xi^* T_{c\xi} = \iota(\langle \xi, c\xi \rangle_A) \in C_0.$$

The second assertion follows from the embedding $C_n \hookrightarrow \mathcal{L}(E^{\otimes n})$ and the fact

$$\|d\| = \sup\{\|\langle \xi, d\xi \rangle_A\| : \xi \in E^{\otimes n}, \|\xi\| = 1\}$$

for $d \in \mathcal{L}(E^{\otimes n})$ positive. $\square$

Lemma 2.7. *Given a positive element $a \in A$ and $\varepsilon > 0$ with $\|a\| > \varepsilon$, there is $\eta \in E$ with $\|\eta\| \leq (\|a\| - \varepsilon)^{-1/2}$ such that $T_\eta^* \iota(a) T_\eta = 1$.*

Proof. Let f be a continuous nonzero real-valued function supported on the interval $[\|a\| - \varepsilon, \|a\|]$ and choose a vector $\zeta \in \pi(f(a))\mathfrak{H}$ such that $\langle \zeta, \pi(a)\zeta \rangle = 1$; we have

$$(\|a\| - \varepsilon)\|\zeta\|^2 \leq \|\langle \zeta, \pi(a)\zeta \rangle\| = 1.$$

Then $\eta = \zeta \otimes 1 \in E$ satisfies the desired conditions. $\square$

It will now follow that $\mathcal{O}_E$ is simple and purely infinite (cf. proof of [Rø, Theorem 2.1]).

Theorem 2.8. *For every nonzero positive element $d \in \mathcal{O}_E$ there is a $z \in \mathcal{O}_E$ so that $z^* dz = 1$. Hence, $\mathcal{O}_E$ is simple and purely infinite.*

Proof. Let $d \in \mathcal{O}_E$ be a nonzero positive element and choose ε so that $0 < \varepsilon < \|P(d)\|/4$. By Proposition 2.4 there is a $\mathbb{T}$ -equivariant isomorphism ψ from $\mathcal{O}_E$ onto a corner of $(\mathcal{O}_E \rtimes_\lambda \mathbb{T}) \rtimes_{\widehat{\lambda}} \mathbb{Z}$ determined by a projection $p \in \mathcal{O}_E \rtimes_\lambda \mathbb{T}$. We now apply Lemma 2.5 to the element $y = \psi(d)$ and the automorphism $\beta = \widehat{\lambda}_1$ (note $\Gamma(\widehat{\lambda}_1) = \mathbb{T}$ by Lemma 2.3). We identify $\mathcal{O}_E$ with the corner determined by p ; note that under this identification $\mathcal{F}_E$ is identified with $p(\mathcal{O}_E \rtimes_\lambda \mathbb{T})p$. There are then positive elements $x, b \in \mathcal{F}_E$ so that

$$\|b\| > \|P(d)\| - \varepsilon, \quad \|x\| \leq 1 \quad \text{and} \quad \|xdx - b\| < \varepsilon.$$

Since $\cup_n C_n$ is dense in $\mathcal{F}_E$ we may assume that $b \in C_n$ for some n . Hence, by Lemma 2.6 there is $\xi \in E^{\otimes n}$ with $\|\xi\| = 1$ such that

$$T_\xi^* b T_\xi \in C_0 \quad \text{and} \quad \|T_\xi^* b T_\xi\| > \|b\| - \varepsilon.$$

Let a denote the unique element of A such that $\iota(a) = T_\xi^* b T_\xi$; then $\|a\| > \|P(d)\| - 2\varepsilon$ and

$$\|T_\xi^* xdx T_\xi - \iota(a)\| = \|T_\xi^* (xdx - b) T_\xi\| < \varepsilon.$$

By Lemma 2.7 there is $\eta \in E$ such that $T_\eta^* \iota(a) T_\eta = 1$ and

$$\|\eta\| \leq (\|a\| - \varepsilon)^{-1/2} < (\|P(d)\| - 3\varepsilon)^{-1/2} < \varepsilon^{-1/2}.$$

It follows that

$$\|T_\eta^* T_\xi^* xdx T_\xi T_\eta - 1\| = \|T_\eta^* (T_\xi^* xdx T_\xi - \iota(a)) T_\eta\| \leq \|T_\xi^* xdx T_\xi - \iota(a)\| (\varepsilon^{-1/2})^2 < 1.$$

Therefore, $c = T_\eta^* T_\xi^* xdx T_\xi T_\eta$ is an invertible positive element and we take $z = x T_\xi T_\eta c^{-1/2}$. $\square$

3. APPLICATIONS AND CONCLUDING REMARKS

We collect some applications of the above theorem and consider certain connections with the theory of reduced (amalgamated) free product C^* -algebras. First we consider criteria under which the Kirchberg-Phillips Theorem applies (see [Kr, Theorem C], [Ph, Corollary 4.2.2]).

Theorem 3.1. *Let A be a separable nuclear unital C^* -algebra which belongs to the bootstrap class to which the UCT applies (see [RS]); let $\pi : A \rightarrow \mathcal{L}(\mathfrak{H})$ be a faithful representation of A on a separable Hilbert space $\mathfrak{H}$ such that $\pi(A) \cap \mathcal{K}(\mathfrak{H}) = \{0\}$ and let E denote the Hilbert A -bimodule $\mathfrak{H} \otimes_{\mathbb{C}} A$. Then $\mathcal{O}_E$ is a unital Kirchberg algebra (simple, purely infinite, separable and nuclear) which belongs to the bootstrap class. Hence, the Kirchberg-Phillips Theorem applies and the isomorphism class of $\mathcal{O}_E$ only depends on $(K_*(A), [1_A])$ and not on the choice of representation π .*

Proof. First note that $\mathcal{O}_E$ is simple and purely infinite by Theorem 2.8. If A is nuclear, then the argument given in the proof of [DS, Theorem 2.1] shows that $\mathcal{O}_E$ must also be nuclear (alternatively, the nuclearity of $\mathcal{O}_E$ follows from the structural results discussed in §1). Hence, $\mathcal{O}_E$ is a unital Kirchberg algebra. Recall that the inclusion $A \hookrightarrow \mathcal{O}_E$ defines a KK -equivalence (see [Pm, Corollary 4.5]) which induces a unit-preserving isomorphism $K_*(A) \cong K_*(\mathcal{O}_E)$. Hence, if A is in the bootstrap class, then $\mathcal{O}_E$ is also. Therefore, the Kirchberg-Phillips Theorem applies and the isomorphism class of $\mathcal{O}_E$ only depends on $(K_*(A), [1_A])$. $\square$

Let X be a second countable compact space, let μ be a nonatomic Borel measure with full support and let

$$\pi : C(X) \rightarrow \mathcal{L}(L^2(X, \mu))$$

be the representation given by multiplication of functions. Then π is faithful and

$$\pi(C(X)) \cap \mathcal{K}(L^2(X, \mu)) = \{0\}.$$

Hence, we may apply the above theorem with $A = C(X)$ and $\mathfrak{H} = L^2(X, \mu)$.

Corollary 3.2. *Let X and μ be as above. Then*

$$E = L^2(X, \mu) \otimes_{\mathbb{C}} C(X)$$

is a Hilbert bimodule over $C(X)$ and $\mathcal{O}_E$ is a unital Kirchberg algebra. The embedding $C(X) \hookrightarrow \mathcal{O}_E$ induces a (unit preserving) KK -equivalence. Hence, the isomorphism class of $\mathcal{O}_E$ only depends on $(K_*(C(X)), [1_{C(X)}])$ (and not on μ); moreover, if X is contractible, then $\mathcal{O}_E \cong \mathcal{O}_{\infty}$.

The following proposition is Theorem 5.6 of [L] (see also [Ka, Theorem 3]); Lance calls this the Kasparov-Stinespring-Gelfand-Naimark-Segal construction.

Proposition 3.3. *Let B and C be C^* -algebras, let F be a Hilbert C -module and let $f : B \rightarrow \mathcal{L}(F)$ be a completely positive map, then there is a Hilbert C -module E_f , a $*$ -homomorphism $\varphi_f : B \rightarrow \mathcal{L}(E_f)$ and an element $v_f \in \mathcal{L}(F, E_f)$ such that $f(b) = v_f^* \varphi_f(b) v_f$ and $\varphi_f(B) v_f F$ is dense in E_f .*

I am grateful to D. Shlyakhtenko for the following observation. Let $\mathcal{T}$ denote the “usual” Toeplitz algebra (i.e. $\mathcal{T}_E$ where E is the 1-dimensional Hilbert bimodule over $\mathbb{C}$) and let g denote the vacuum state on $\mathcal{T}$.

Proposition 3.4. *Let A be a separable unital C^* -algebra and let $\pi : A \rightarrow \mathcal{L}(\mathfrak{H})$ be a faithful representation of A on a separable Hilbert space $\mathfrak{H}$ such that π has a cyclic vector $\xi \in \mathfrak{H}$. Let f denote the vector state $\langle \xi, \pi(\cdot) \xi \rangle$ and let $\tilde{f}$ denote the corresponding completely positive map from A to $\mathcal{L}(A)$ (given by $\tilde{f}(a) = f(a)1$). Then $E = E_{\tilde{f}} \cong \mathfrak{H} \otimes A$ and $\mathcal{T}_E$ may be realized as a reduced free product (see [A, V]):*

$$(\mathcal{T}_E, h) \cong (A, f) * (\mathcal{T}, g)$$

for some state h on $\mathcal{T}_E$.

Proof. This follows from [Sh, Theorem 2.3, Corollary 2.5]. $\square$

As a result of this observation part (at least) of Corollary 3.2 follows from the existing literature on reduced free products. The simplicity follows from a theorem of Dykema [Dy, Theorem 2]. Criteria for when reduced free products are purely infinite have been found by Choda, Dykema and Rørdam in a series of papers [DR1, DR2, DC]; but none seem to apply generally to the case considered in the corollary.

A theorem of Speicher (see [Sp]) on reduced amalgamated free products (see [V, §5]) and Toeplitz algebras associated to Hilbert bimodules yields a curious stability property of the algebras we have been considering. The following is the version given in [BDS, Theorem 2.4].

Proposition 3.5. *Suppose that E_1 and E_2 are full Hilbert bimodules over the C^* -algebra A . Then*

$$\mathcal{T}_{E_1 \oplus E_2} = \mathcal{T}_{E_1} *_A \mathcal{T}_{E_2}.$$

We obtain the following corollary.

Corollary 3.6. *Let A be a separable nuclear unital C^* -algebra which belongs to the bootstrap class to which the UCT applies (see [RS]) and let $\pi : A \rightarrow \mathcal{L}(\mathfrak{H})$ be a faithful representation of A on a separable Hilbert space $\mathfrak{H}$ such that $\pi(A) \cap \mathcal{K}(\mathfrak{H}) = \{0\}$. Let E be the Hilbert bimodule $\mathfrak{H} \otimes_{\mathbb{C}} A$. Then*

$$\mathcal{O}_E \cong \mathcal{O}_E *_A \mathcal{O}_E.$$

Proof. Observe that $E \oplus E = (\mathfrak{H} \oplus \mathfrak{H}) \otimes_{\mathbb{C}} A$. Since $\pi \oplus \pi : A \rightarrow \mathcal{L}(\mathfrak{H} \oplus \mathfrak{H})$ is a faithful representation and $(\pi \oplus \pi)(A) \cap \mathcal{K}(\mathfrak{H} \oplus \mathfrak{H}) = \{0\}$, the result follows from Theorem 3.1 and the above proposition. $\square$

REFERENCES

- [A] D. Avitzour, *Free products of C^* -algebras*, Trans. Amer. Math. Soc. **271** (1982) 423–435.
- [BDS] N. Brown, K. Dykema and D. Shlyakhtenko, *Topological Entropy of Free Product Automorphisms*, preprint.
- [Co] A. Connes, *Une classification des facteurs de type III*, Ann. Sci. École Norm. Sup. **6** (1973) 133–252.
- [Cu] J. Cuntz, *Simple C^* -algebras generated by isometries*, Commun. Math. Phys. **57** (1977) 173–185.
- [CK] J. Cuntz and W. Krieger, *A class of C^* -algebras and topological Markov chains*, Invent. Math. **56** (1980) 251–268.
- [De] V. Deaconu, *Generalized solenoids and C^* -algebras*, Pacific J. Math. **190** (1999), 247–260.
- [DPW] Doplicher, C. Pinzari, and R. Zuccante, *The C^* -algebra of a Hilbert bimodule*, Boll. Unione Mat. Ital. Sez. B Artic. Ric. Mat. (8) **1** (1998) 263–281.
- [Dy] K. Dykema, *Simplicity and the stable rank of some free product C^* -algebras*, Trans. Amer. Math. Soc. **351** (1999) 1–40.
- [DC] K. Dykema and M. Choda, *Purely infinite, simple C^* -algebras arising from free product constructions, III*, Proc. Amer. Math. Soc. **128** (2000) 3269–3273.
- [DR1] K. Dykema and M. Rørdam, *Purely infinite, simple C^* -algebras arising from free product constructions*, Canad. J. Math. **50** (1998) 323–341.
- [DR2] K. Dykema and M. Rørdam, *Purely infinite, simple C^* -algebras arising from free product constructions, II*, preprint.
- [DS] K. Dykema and D. Shlyakhtenko, *Exactness of Cuntz-Pimsner C^* -algebras*, Proc. Edinburgh Math. Soc., to appear.
- [E] George A. Elliott, *Some simple C^* -algebras constructed as crossed products with discrete outer automorphism groups*, Publ. Res. Inst. Math. Sci. **16** (1980) 299–311.
- [FR] N. J. Fowler and I. Raeburn, *The Toeplitz algebra of a Hilbert bimodule*, Indiana Univ. Math. J. **48** (1999) 155–181.
- [FRM] N. J. Fowler, I. Raeburn and P. S. Muhly, *Representations of Cuntz-Pimsner C^* -algebras*, preprint.
- [KPW1] T. Kajiwara, C. Pinzari and Y. Watatani, *Ideal structure and simplicity of the C^* -algebras generated by Hilbert bimodules*, J. Funct. Anal. **159** (1998) 295–322.

- [KPW2] T. Kajiwara, C. Pinzari and Y. Watatani, *Hilbert C^* -bimodules and countably generated Cuntz-Krieger algebras*, J. Operator Theory **45** (2001) 3–18.
- [Ka] Kasparov, G. G. *Hilbert C^* -modules: theorems of Stinespring and Voiculescu*, J. Operator Theory **4** (1980) 133–150.
- [Kr] E. Kirchberg, *The classification of purely infinite C^* -algebras using Kasparov’s theory*, Fields Institute Communications, to appear.
- [Ks] A. Kishimoto, *Outer automorphisms and reduced crossed products of simple C^* -algebras*, Comm. Math. Phys. **81** (1981) 429–435.
- [L] E. C. Lance, *Hilbert C^* -modules. A toolkit for operator algebraists*, London Math. Soc. Lecture Note Series, **210**, Cambridge University Press, Cambridge, 1995.
- [MS] P. S. Muhly and B. Solel, *On the simplicity of some Cuntz-Pimsner algebras*, Math. Scand. **83** (1998) 53–73.
- [O] D. Olesen, *Inner $*$ -automorphisms of simple C^* -algebras*, Comm. Math. Phys. **44** (1975) 175–190.
- [OP1] D. Olesen and G. K. Pedersen, *Applications of the Connes Spectrum to C^* -dynamical Systems*, J. Funct. Anal. **30** (1978) 179–197.
- [OP2] D. Olesen and G. K. Pedersen, *Applications of the Connes Spectrum to C^* -dynamical Systems, II*, J. Funct. Anal. **36** (1980) 18–32.
- [OP3] D. Olesen and G. K. Pedersen, *Applications of the Connes Spectrum to C^* -dynamical Systems, III*, J. Funct. Anal. **45** (1982) 357–390.
- [Pa] W. L. Paschke, *The crossed product of a C^* -algebra by an endomorphism*, Proc. Amer. Math. Soc. **80** (1980) 113–118.
- [Pd] G. K. Pedersen, *C^* -algebras and their automorphism groups*, Academic Press, London, 1979.
- [Ph] N. C. Phillips, *A classification theorem for nuclear purely infinite simple C^* -algebras*, Doc. Math. **5** (2000) 49–114.
- [Pm] M. Pimsner, *A class of C^* -algebras generalizing both Cuntz-Krieger algebras and crossed products by $\mathbb{Z}$* , in “Free probability theory” (D. Voiculescu, Ed.), 189–212, Fields Institute Communications **12**, Amer. Math. Soc., Providence, 1997.
- [PV] M. Pimsner and D. Voiculescu, *Exact sequences for K -groups and Ext-groups of certain cross-product C^* -algebras*, J. Operator Theory **4** (1980) 93–118.
- [Pn] C. Pinzari, *The ideal structure of Cuntz-Krieger-Pimsner algebras and Cuntz-Krieger algebras over infinite matrices*, in “Operator Algebras and Quantum Field Theory (Accademia Nazionale dei Lincei, Roma, July 1-6 1996)” (S. Doplicher et al, Eds.), 136–150, International Press 1997.
- [Ri] M. A. Rieffel, *Induced Representations of C^* -algebras*, Advances Math. **13** (1974) 176–257.
- [Rø] M. Rørdam, *Classification of certain infinite simple C^* -algebras, III*, in “Operator algebras and their applications” (P. Fillmore and J. Mingo, Eds.), 257–283, Fields Institute Communications **13**, Amer. Math. Soc., Providence, 1997.
- [Ro] J. Rosenberg, *Appendix to: “Crossed products of UHF algebras by product type actions,” by O. Bratteli*, Duke Math. J. **46** (1979) 25–26.
- [RS] J. Rosenberg and C. Schochet, *The Künneth theorem and the universal coefficient theorem for Kasparov’s generalized K -functor*, Duke Math. J. **55** (1987) 431–474.
- [Sc1] J. Schweizer, *Crossed products by C^* -correspondences and Cuntz-Pimsner algebras*, in “ C^* -algebras, Münster, 1999” (J. Cuntz and S. Echterhoff, Eds.), 203–226, Springer Verlag, Berlin, 2000.
- [Sc2] J. Schweizer, *Dilations of C^* -correspondences and the simplicity of Cuntz-Pimsner algebras*, J. Funct. Anal. **180** (2001) 404–425.
- [Sh] D. Shlyakhtenko, *Some applications of freeness with amalgamation*, J. Reine Angew. Math. **500** (1998), 191–212.
- [Sp] R. Speicher, *Combinatorial theory of the free product with amalgamation and operator-valued free probability theory*, Mem. Amer. Math. Soc. **132**, no. 627 (1998).
- [T] H. Takai, *On a duality for crossed products of C^* -algebras*, J. Funct. Anal. **19** (1975), 25–39.
- [V] D. Voiculescu, *Symmetries of some reduced free product C^* -algebras*, in “Operator algebras and their connections with topology and ergodic theory, Buşteni, 1983,” (H. Araki et al, Eds.) 556–588, Lecture Notes in Math., 1132, Springer, Berlin, 1985.
- [Z] J. Zacharias, *Quasi-free automorphisms of Cuntz-Krieger-Pimsner algebras*, in “ C^* -algebras, Münster 1999.” (J. Cuntz and S. Echterhoff, Eds.), 262–272, Springer Verlag, Berlin, 2000.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF NEVADA, RENO NV 89557, USA

E-mail address: alex@unr.edu